



ORIGINAL RESEARCH ARTICLE

Open Access



Investigating effect of *Helicobacter pylori* treatment on improvement of non-alcoholic fatty liver parameters: a randomized trial

Amir Mohammad Kazemifar¹, Ali Akbar Shafikhani², Elaheh HajiNoormohammadi³, Zahra Azarion³ and Aliakbar Hajiaghamohammadi^{4*}

Abstract

Background and objective: The correlation between the eradication of *Helicobacter pylori* (HP) and non-alcoholic fatty liver disease (NAFLD) is a controversial one. The aim of this study is to investigate the effect of *Helicobacter pylori* treatment on liver function tests and lipid profiles and to compare its effects with exercise therapy and diet alone.

Method: This was a double-blind randomized clinical trial conducted at Qazvin University of Medical Sciences. One hundred patients with NAFLD having a positive test for urea breath testing were randomly assigned into one of the intervention and comparison groups. The diet and physical activity program were given to two groups of patients for 8 weeks, three sessions per week. In addition to the above therapy, the patients in the intervention group also received HP treatment for 2 weeks. The data corresponding to anthropometric and clinical features before and after the intervention were collected in both groups and compared using appropriate statistical methods.

Results: After the treatment interventions, the variables of weight, BMI, blood glucose, triglyceride, AST, ALT, total cholesterol, and LDL-C were significantly decreased in both groups ($p < .05$). In the between-group comparisons, only ALT was significantly lower in the intervention group (HP) ($p < .05$).

Conclusion: Findings of this study showed that the eradication of *Helicobacter pylori* could significantly improve the ALT index, but it had no additional effect on changes in metabolic indicators.

Trial registration: Registration number: IRCT2015042020951N2

Name of trial registry: The study of treatment of *Helicobacter pylori* in improvement of non-alcoholic fatty liver disease
The date of registration: 2015-05-12

Where the full trial protocol can be accessed: <https://fa.irct.ir/trial/18489>

Keywords: Non-alcoholic fatty liver disease, *Helicobacter pylori*, Alanine transaminase, Aspartate aminotransferases

Background

Non-alcoholic fatty liver disease (NAFLD) is a type of liver injury caused by metabolic stress and hereditary sensitivity. Generally, this disease is considered to be the liver manifestation of the metabolic syndrome [1, 2]. NAFLD includes non-alcoholic fatty liver (NAFL), non-alcoholic steatohepatitis (NASH), and non-alcoholic steatohepatitis associated

with hepatic cirrhosis and hepatocellular carcinoma. The incidence of NAFLD complications is rapidly increasing in the world [2, 3]; therefore, the economic and clinical burden of this disease is significant. Recently, except for the known risk factors such as obesity, type 2 diabetes, hypertension, and dyslipidemia, the role of the pathogenesis of *Helicobacter pylori* (HP) is also discussed in the disease and may be associated with NAFLD [4, 5].

HP is a Gram-negative bacterium that is selectively placed in the gastric mucosa. This bacterium is the cause of most cases of gastric ulcers and leads to development of some

* Correspondence: ahmohammadi@qums.ac.ir

⁴Gastroenterologist Metabolic Diseases Research Center, Qazvin University of Medical Sciences, Qazvin, Iran

Full list of author information is available at the end of the article

body weight could result in the sustained biochemical improvement. Given the limitations expressed in the diagnosis of the disease, for future studies, it is recommended to use biopsy and magnetic resonance elastography (MRE) to diagnose and MRE to follow up.

Abbreviations

ALT: Alanine aminotransferase; AST: Aspartate aminotransferase; HDL: High-density lipoprotein; HP: *Helicobacter pylori*; LDL: Low-density lipoprotein; NAFL: Non-alcoholic fatty liver; NAFLD: Non-alcoholic fatty liver disease; NASH: Non-alcoholic steatohepatitis

Acknowledgements

The researchers would like to acknowledge and appreciate the contributions of all clinical residents of the Qazvin University of Medical Sciences who helped in conducting the present study.

Authors' contributions

All authors read and approved the final manuscript. AK did the design, data gathering, and manuscript preparation. AS did the data gathering, analysis, and manuscript preparation. EH did the data gathering and manuscript preparation. ZA did the data gathering and manuscript preparation. AH did the design, data gathering, analysis, and manuscript preparation.

Funding

Deputy for research of Qazvin University of Medical Sciences.

Availability of data and materials

Data and materials were available for study.

Ethics approval and consent to participate

This study was approved by the Ethics Committee of Qazvin University of Medical Sciences and the participation of individuals was subject to a written consent (Ethics committee reference number: 28/20/9669).

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Clinical Toxicology, Metabolic Diseases Research Center, Qazvin University of Medical Sciences, Qazvin, Iran. ²Department of Occupational Health Engineering, Shahid Beheshti University of Medical Sciences, Tehran, Iran. ³Department of Internal Medicine, Metabolic Disease Research Center, Qazvin University of Medical Sciences, Qazvin, Iran. ⁴Gastroenterologist Metabolic Diseases Research Center, Qazvin University of Medical Sciences, Qazvin, Iran.

Received: 17 July 2019 Accepted: 15 August 2019

Published online: 10 October 2019

References

- Benedict M, Zhang X (2017) Non-alcoholic fatty liver disease: an expanded review. *World J Hepatol* 9(16):715
- Neuschwander-Tetri BA (2017) Non-alcoholic fatty liver disease. *BMC Med* 15(1):45
- Engin AB, Engin A (2017) Obesity and lipotoxicity: Springer
- Cai O, Huang Z, Li M, Zhang C, Xi F, Tan S (2018) Association between *Helicobacter pylori* infection and nonalcoholic fatty liver disease: a single-center clinical study. *Gastroenterol Res Pract* 2018:8040262 PubMed Central PMCID: PMC5828541. Epub 2018/03/13. eng
- Kang S, Kim D (2016) PTH-112 association between *Helicobacter pylori* infection and nonalcoholic fatty liver disease in the United States. *BMJ Publishing Group* 65: A273–A274
- Malfertheiner P, Megraud F, O'Morain CA, Gisbert JP, Kuipers EJ, Axon AT et al (2017) Management of *Helicobacter pylori* infection-the Maastricht V/Florence consensus report. *Gut* 66(1):6–30 PubMed PMID: 27707777. Epub 2016/11/02. eng
- Buruco C, Axon A (2017) Epidemiology of *Helicobacter pylori* infection. *Helicobacter* 22(Suppl 1):e12403 PubMed PMID: 28891138. Epub 2017/09/12. eng
- Eusebi LH, Zagari RM, Bazzoli F (2014) Epidemiology of *Helicobacter pylori* infection. *Helicobacter* 19(Suppl 1):1–5 PubMed PMID: 25167938. Epub 2014/08/30. eng
- Leja M, Axon A, Brenner H (2016) Epidemiology of *Helicobacter pylori* infection. *Helicobacter* 21(Suppl 1):3–7 PubMed PMID: 27531531. Epub 2016/08/18. eng
- Franceschi F, Gasbarrini A, Polyzos SA, Kountouras J (2015) Extragastric diseases and *Helicobacter pylori*. *Helicobacter* 20(Suppl 1):40–46 PubMed PMID: 26372824. Epub 2015/09/16. eng
- Nasif WA, Mukhtar MH, Nour Eldein MM, Ashgar SS (2016) Oxidative DNA damage and oxidized low density lipoprotein in Type II diabetes mellitus among patients with *Helicobacter pylori* infection. *Diabetol Metab Syndr* 8: 34 PubMed PMID: 27148410. eng
- Taylor NS, Fox JG, Yan L (1995) In-vitro hepatotoxic factor in *Helicobacter hepaticus*, *H. pylori* and other *Helicobacter* species. *J Med Microbiol* 42(1): 48–52 PubMed PMID: 7739025. Epub 1995/01/01. eng
- Takuma Y (2011) *Helicobacter pylori* infection and liver diseases. *Gan To Kagaku Ryoho* 38(3):362–364 PubMed PMID: 21403438. Epub 2011/03/16. jpn
- Jamali R, Mofid A, Vahedi H, Farzaneh R, Dowlatabadi S (2013) The effect of *Helicobacter pylori* eradication on liver fat content in subjects with non-alcoholic fatty liver disease: a randomized open-label clinical trial. *Hepat Mon* 13(12):e14679 PubMed PMID: 24358044. eng
- Hashemi MR, Rahnavardi M, Bikieli B, Dehghani ZM (2006) *H. pylori* infection among 1000 southern Iranian dyspeptic patients. *World J Gastroenterol* 12(34):5479–5482 PubMed PMID: 17006984. Epub 09/14. eng
- Lankarani KB, Ghaffarpasand F, Mahmoodi M, Lotfi M, Zamiri N, Heydari ST et al (2013) Non alcoholic fatty liver disease in southern Iran: a population based study. *Hepat Mon* 13(5):e9248 PubMed PMID: 23922564. eng
- Ambade VN, Sharma YV, Somani BL (1998) Methods for estimation of blood glucose : a comparative evaluation. *Med J Armed Forces India* 54(2):131 PubMed PMID: 28775446. PubMed Central PMCID: PMC5531325. Epub 1998/04/01. eng
- Jia KK, Zhang J (2010) Evaluation of five routine glucose methods on an Olympus AU5400 analyzer using the CDC hexokinase reference method. *Clin Chem Lab Med* 48(3):361–364 PubMed PMID: 20170396. Epub 2010/02/23. eng
- Knopfholz J, Disserol CC, Pierin AJ, Schirr FL, Streisky L, Takito LL et al (2014) Validation of the friedewald formula in patients with metabolic syndrome. *Cholesterol* 2014:261878 PubMed PMID: 24672715. PubMed Central PMCID: PMC3941209. Epub 2014/03/29. eng
- Jamali R, Khonsari M, Merat S, Khoshnia M, Jafari E, Bahram Kalhori A et al (2008) Persistent alanine aminotransferase elevation among the general Iranian population: prevalence and causes. *World J Gastroenterol* 14(18):2867–2871 PubMed PMID: 18473412. PubMed Central PMCID: PMC2710729. Epub 2008/05/14. eng
- Polyzos SA, Nikolopoulos P, Stogianni A, Romiopoulou I, Katsinelos P, Kountouras J (2014) Effect of *Helicobacter pylori* eradication on hepatic steatosis, NAFLD fibrosis score and HSENLI in patients with nonalcoholic steatohepatitis: a MR imaging-based pilot open-label study. *Arg Gastroenterol* 51(3):261–268 PubMed PMID: 25296089. Epub 2014/10/09. eng
- Okushin K, Takahashi Y, Yamamichi N, Shimamoto T, Enooku K, Fujinaga H et al (2015) *Helicobacter pylori* infection is not associated with fatty liver disease including non-alcoholic fatty liver disease: a large-scale cross-sectional study in Japan. *BMC Gastroenterol* 15:25 PubMed PMID: 25880912. PubMed Central PMCID: PMC4349671. Epub 2015/04/17. eng
- Della Pepa G, Vetrani C, Lombardi G, Bozzetto L, Annuzzi G, Rivellese AA (2017) Isocaloric dietary changes and non-alcoholic fatty liver disease in high cardiometabolic risk individuals. *Nutrients* 9(10):1065
- Slentz CA, Bateman LA, Willis LH, Shields AT, Tanner CJ, Piner LW et al (2011) Effects of aerobic vs. resistance training on visceral and liver fat stores, liver enzymes, and insulin resistance by HOMA in overweight adults from STRRIDE AT/RT. *Am J Physiol Endocrinol Metab* 301(5):E1033–E1039 PubMed PMID: 21846904. PubMed Central PMCID: PMC3214001. Epub 2011/08/19. eng
- Park SK, Park JH, Kwon YC, Kim HS, Yoon MS, Park HT (2003) The effect of combined aerobic and resistance exercise training on abdominal fat in obese middle-aged women. *J Physiol Anthropol Appl Hum Sci* 22(3):129–135 PubMed PMID: 12808225. Epub 2003/06/17. eng
- Sigal RJ, Kenny GP, Boule NG, Wells GA, Prud'homme D, Fortier M et al (2007) Effects of aerobic training, resistance training, or both on glycemic control in type 2 diabetes: a randomized trial. *Ann Intern Med* 147(6):357–369 PubMed PMID: 17876019. Epub 2007/09/19. eng

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.